

JNK1/2/3 (P184) polyclonal antibody

Catalog: BS1544

Host: Rabbit

Reactivity: Human, Mouse, Rat

Background:

The human JNK1 (PRKM8, SAPK1, MAPK8) gene maps to chromosome 10q11.22 and shares 83% amino acid identity with JNK2. JNK1 is necessary for normal activation and differentiation of CD4 helper T (TH) cells into TH1 and TH2 effector cells. Capsaicin activates JNK1 and p38 in Ras-transformed human breast epithelial cells. Nitrogen oxides (NOx) upregulate JNK1 in addition to c-Fos, c-Jun and other signaling kinases, including MEKK1 and p38. JNK3 (MK10, MAPK10, PRKM10) is activated by pro-inflammatory cytokines and environmental stress by phosphorylating transcription factors such as c-Jun and ATF2. This is important for AP-1 transcriptional activity regulation. JNK3 is crucial for neuronal apoptosis (stress-induced).

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 46,54 kDa

Swiss-Prot:

P45983/P45984/P53779

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

JNK1/2/3 (T178) polyclonal antibody detects endogenous levels of JNK1/2/3 protein.

DATA:




Western blot (WB) analysis of JNK1/2/3 (P184) polyclonal antibody at 1:500 dilution

Lane1:HEK293T whole cell lysate


Lane2:Raw264.7 whole cell lysate

Lane3:H9C2 whole cell lysate



BS1544
Lot AA54131

Immunohistochemistry (IHC) analyzes of JNK1/2/3 (P184) pAb in paraffin-embedded human breast carcinoma tissue at 1:50. showing cytoplasmic and nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



BS1544
Lot AA54131

Immunohistochemistry (IHC) analyzes of JNK1/2/3 (P184) pAb in paraffin-embedded human kidney carcinoma tissue at 1:50. showing cytoplasmic and nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.


Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



IgG-biotin followed by avidin-peroxidase.



BS1544
Lot AA54131

Immunohistochemistry (IHC) analyzes of JNK1/2/3 (P184) pAb in paraffin-embedded human rectum carcinoma tissue at 1:50. showing cytoplasmic and nucleus staining. Negative control (the right) Using PBS

instead of primary antibody, secondary antibody is Goat Anti-Rabbit

IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151