

PAK1/2/3 (E417) polyclonal antibody

Catalog: BS1570

Host: Rabbit

Reactivity: Human,Rat,Mouse

BackGround:

Three isoforms of serine/threonine kinases, designated α PAK p68, β PAK p65 and γ PAK p62, have been shown to exhibit a high degree of sequence homology with the *S. cerevisiae* kinase Ste 20, involved in pheromone signaling. The α , β and γ PAK isoforms complex specifically with Rac1 and Cdc42 in their active GTP-bound state, inhibiting their intrinsic GTPase activity leading to their autophosphorylation. There are eight sites of autophosphorylation on γ PAK, including Ser 19, Ser 141 and Thr 402, and phosphorylation of Ser 141 and Thr 402 is correlated with γ PAK activation. Once phosphorylated and their affinity for Rac/Cdc42 reduced, the PAK isoforms disassociate from the complex to seek downstream substrates.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 61 kDa

Swiss-Prot:

Q13153/Q13177/O75914

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:1000~1:2000

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

PAK1/2/3 (E417) polyclonal antibody detects endogenous levels of total PAK protein.

DATA:



Western blot (WB) analysis of PAK1/2/3 (E417) polyclonal antibody at 1:1000 dilution


Lane1:SSGC7901 whole cell lysate(40ug)

Lane2:U-87MG whole cell lysate(40ug)

Lane3:THP-1 whole cell lysate(40ug)

Lane4:RAW264.7 whole cell lysate(40ug)

Lane5:C6 whole cell lysate(40ug)



Immunofluorescence analysis of U-87MG cells using PAK1/2/3 (E417) antibody at dilution of 1:50.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151