

VDR (N204) polyclonal antibody

Catalog: BS1774

Host: Ral

Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

The vitamin D receptor (VDR), also known as the calcitriol receptor, and also known as NR111 (nuclear receptor subfamily 1, group I, member 1), is a member of the nuclear receptor family of transcription factors. Upon activation by vitamin D, the VDR forms a heterodimer with the retinoid-X receptor and binds to hormone response elements on DNA resulting in expression or trans-repression of specific gene products. It is an intracellular hormone receptor that specifically binds 1,25(OH)2D3 and mediates its effects. Downstream targets of this nuclear hormone receptor are principally involved in mineral metabolism though the receptor regulates a variety of other metabolic pathways, such as those involved in the immune response and cancer.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 43, 55 kDa

Swiss-Prot:

P11473

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.


Specificity:

VDR (N204) polyclonal antibody detects endogenous levels of VDR protein


Bioworld Technology, Inc.

Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841


DATA:



Western blot (WB) analysis of VDR (N204) pAb at 1:500 dilution Lane1:K562 whole cell lysate(40ug) Lane2:HCT116 whole cell lysate(40ug) Lane3:3T3-L1 whole cell lysate(40ug) Lane4:The Kidney tissue lysate of Rat(40ug)



Immunohistochemistry (IHC) analyzes of VDR (N204) pAb in paraffin-embedded human tonsil carcinoma tissue at 1:50.



Western blot (WB) analysis of VDR pAb at 1:1000 dilution Lane1:HCT116 whole cell lysate(40ug) Lane2:HEK293T whole cell lysate(40ug) Lane3:A549 whole cell lysate(40ug) Lane4:CT26 whole cell lysate(40ug) Lane5:PC12 whole cell lysate(40ug)

Note:

Bioworld technology, co. Ltd.Add:No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.Email:info@biogot.comTel:0086-025-68037686Fax:0086-025-68035151



Bioworld Technology, Inc.

For research use only, not for use in diagnostic procedure.

Bioworld	Technol	loqy,	Inc.
----------	---------	-------	------

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 Fax: 6122933841

Bioworld technology, co. Ltd.

 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151