

## IKK $\gamma$ (D25) polyclonal antibody

Catalog: BS1816

Host: Rabbit

Reactivity: Human, Mouse, Rat

### Background:

Activation of NF $\kappa$ B requires that I $\kappa$ B be phosphorylated on specific serine residues, which results in targeted degradation of I $\kappa$ B. I $\kappa$ B kinase  $\alpha$  (IKK $\alpha$ ), previously designated CHUK, interacts with I $\kappa$ B- $\alpha$  and specifically phosphorylates I $\kappa$ B $\alpha$  on Serine 32 and 36, the sites that trigger its degradation. IKK $\alpha$  appears to be critical for NF $\kappa$ B activation in response to proinflammatory cytokines. Phosphorylation of I $\kappa$ B by IKK $\alpha$  is stimulated by the NF $\kappa$ B inducing kinase (NIK), which itself is a central regulator for NF $\kappa$ B activation in response to TNF and IL-1. The functional IKK complex contains three subunits, IKK $\alpha$ , IKK $\beta$  and IKK $\gamma$  (also designated NEMO), and each appear to make essential contributions to I $\kappa$ B phosphorylation.

### Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

### Molecular Weight:

~ 48 kDa

### Swiss-Prot:

Q9Y6K9

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


### Storage&Stability:

Store at 4  $^{\circ}$ C short term. Aliquot and store at -20  $^{\circ}$ C long term. Avoid freeze-thaw cycles.

### Specificity:

IKK $\gamma$  (D25) polyclonal antibody detects endogenous levels of IKK $\gamma$  protein.

### DATA:




Western blot (WB) analysis of IKK $\gamma$  (D25) polyclonal antibody at 1:5000 dilution

Lane1: Jurkat whole cell lysate (30ug)

Lane2: RAW264.7 whole cell lysate (30ug)

Lane3: H9C2 whole cell lysate (30ug)

Lane4: NIH-3T3 whole cell lysate (30ug)



Immunohistochemistry (IHC) analysis of IKK $\gamma$  (D25) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

### Note:

For research use only, not for use in diagnostic procedure.

### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151