

TFIIE- β (K183) polyclonal antibody

Catalog: BS1974

Host: Rabbit

Reactivity: Human, Mouse

BackGround:

The general transcription factor TFIIE consists of two subunits, α and β , and plays essential roles in transcription. Structure-function studies indicate that TFIIE has three-winged helix (WH) motifs, with one in TFIIE α and two in TFIIE β . By binding to the clamp region of RNA polymerase II, TFIIE α -WH promotes the conformational change that transforms the promoter-bound inactive preinitiation complex to the active complex.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 33 kDa

Swiss-Prot:

P29084

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

TFIIE- β (K183) pAb detects endogenous levels of TFIIE- β protein.

DATA:



Western blot (WB) analysis of TFIIE- β (K183) pAb at 1:500 dilution

Lane1: SGC7901 whole cell lysate (40ug)

Lane2: HCT116 whole cell lysate (40ug)

Lane3: The Testis tissue lysate of Mouse (40ug)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151