

PERK (R87) polyclonal antibody

Catalog: BS2156

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

An interferon-inducible, RNA-dependent protein serine /threonine kinase (PKR) has been described. PKR in earlier literature is variously known as DAI, dsJ, PI kinase, p65, p67 or TIK for the mouse kinase; and p68 or p69 for the human kinase. The PKR kinase substrate is the α subunit of protein synthesis initiation factor eIF-2. Phosphorylation of eIF-2 α on serine-51 results in inhibition of translation. The serine /threonine kinase catalytic domains map to the carboxy terminal half of the protein while the RNA-binding domains are located in the amino terminal region. PERK is a type I transmembrane protein located in the endoplasmic reticulum (ER) that contains a kinase domain similar to the kinase domain of PKR. PERK is activated in response to ER stress and phosphorylates eIF-2 α , thus inhibiting the translation of mRNA.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 140 kDa

Swiss-Prot:

Q9NZJ5

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200

Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

PERK (R87) polyclonal antibody detects endogenous

levels of PERK protein.

DATA:



Western blot (WB) analysis of PERK (R87) pAb at 1:1000 dilution


Lane1:HCT116 whole cell lysate(20ug)

Lane2:MCF-7 whole cell lysate(20ug)


Lane3:A375 whole cell lysate(20ug)

Lane4:CT26 whole cell lysate(40ug)

Lane5:PMVEC whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of PERK (R87) pAb in paraffin-embedded human colon carcinoma tissue at 1:50. showing cytoplasmic and nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



Immunohistochemistry (IHC) analyzes of PERK (R87) pAb in paraffin-embedded human liver carcinoma tissue at 1:50. showing cytoplasmic and nucleus staining.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology, Inc.

mic and nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151