

# PRODUCT DATA SHEET

Bioworld Technology,Inc.

# TG1019 (R251) polyclonal antibody

Catalog: BS3237 Host: Rabbit Reactivity: Human, Mouse, Rat

## **BackGround:**

TG1019 is a G protein-coupled receptor that functions as a chemoattractant for eosinophils and neutrophils. TG1019 is expressed in various tissues except the brain, with the highest expression in liver, kidney, peripheral leukocyte, lung and spleen. TG1019 interacts with the protein OXE to form TG1019/OXE, a Ga i/o protein-coupled receptor. Signals from TG1019 are transduced via a Ga i/o protein to PLC/calcium, MEK/ERK and PI3K/Akt pathways. Signal transduction from TG1019 following stimulation with 5-oxo-6E,8Z,11Z,14Z eicosatetraenoic acid (5-oxo-ETE) induces intracellular calcium mobilization and rapid activation of MEK/ERK and PI3K/Akt pathways. TG1019 also may play a role in downregulating cAMP production.

### **Product:**

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

## **Molecular Weight:**

~ 46 kDa

#### **Swiss-Prot:**

Q8TDS5

## **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

#### **Applications:**

WB: 1:500~1:1000 IHC: 1:50~1:200 IF: 1:50~1:200


## Storage&Stability:

Store at  $4 \,\mathrm{C}$  short term. Aliquot and store at  $-20 \,\mathrm{C}$  long term. Avoid freeze-thaw cycles.

## **Specificity:**

TG1019 (R251) polyclonal antibody detects endogenous levels of TG1019 protein.

#### **DATA:**




Western blot (WB) analysis of TG1019 (R251) pAb at 1:500 dilution

Lane1:3T3-L1 whole cell lysate(40ug)

Lane2:AML-12 whole cell lysate(40ug)

Lane3:H1792 whole cell lysate(40ug)

Lane4:L02 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of TG1019 (R251) pAb in paraffin-embedded human liver carcinoma tissue at 1:50.showing cytoplasmic staining. Negative control (the right)Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

#### Note

For research use only, not for use in diagnostic procedure.

## Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <a href="mailto:info@biogot.com">info@biogot.com</a>
Tel: 0086-025-68037686
Fax: 0086-025-68035151