

TRAF6 (H154) polyclonal antibody

Catalog: BS3684

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Tumor necrosis factor receptor-associated factor 6 (TRAF6) regulates adaptive immunity, innate immunity and bone metabolism. TRAF6 is a ubiquitin (Ub) ligase that mediates the activation of protein kinases, such as transforming growth factor beta-activated kinase (TAK1) and I κ B kinase (IKK), by catalyzing the formation of a unique polyubiquitin chain linked through Lys 63 of Ub. TRAF6 is essential for activating NF κ B signaling pathway in response to interleukin-1 and Toll-like receptor ligands. The coiled-coil domain of TRAF6 is essential for its auto-ubiquitination and activating NF κ B signaling pathway. TRAF6 interacts with various protein kinases including IRAK1/IRAK, SRC and PKC ζ , which provides a link between distinct signaling pathways.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 58 kDa

Swiss-Prot:

Q9Y4K3

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

TRAF6 (H154) polyclonal antibody detects endogenous levels of TRAF6 protein.

DATA:




Western blot (WB) analysis of TRAF6 (H154) polyclonal antibody at 1:500 dilution

Lane1:MCF-7 cell lysate

Lane2:Mouse kidney tissue lysate


Lane3:Rat kidney tissue lysate

Lane4:HEK293T cell lysate



BS3684
Lot C136131

Immunohistochemistry (IHC) analyzes of TRAF6 (H154) pAb in paraffin-embedded human colon carcinoma tissue at 1:50, showing cytoplasmic and nuclear staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



BS3684
Lot C136131

Immunohistochemistry (IHC) analyzes of TRAF6 (H154) pAb in paraffin-embedded human tonsil carcinoma tissue at 1:50, showing cytoplasmic and nuclear staining. Negative control (the right) Using PBS instead

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET


Bioworld Technology, Inc.

of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.



BS3684
Lot C136131

Immunohistochemistry (IHC) analyzes of TRAF6 (H154) pAb in paraffin-embedded human liver carcinoma tissue at 1:50, showing cytoplasmic and nuclear staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151