

SMC 1 (phospho-S957) polyclonal antibody

Catalog: BS4174

Host: Rabbit

Reactivity: Human

BackGround:

The SMC (structural maintenance of chromosomes) family of proteins form heterodimeric complexes that modulate sister chromatid cohesion and chromosome condensation for mitosis. The two distinct classes of SMC protein complexes are comprised of SMC1 (also designated SB1.8) with SMC3 (also designated HCAP for human chromosome-associated protein, and Bamacan), and SMC2 (also designated hCAP-E) with SMC4 (also designated hCAP-C). The SMC1/SMC3 complex is required for metaphase progression in mitotic cells and functions independently of the SMC2/SMC4 complex during the cell cycle. SMC1 is ubiquitously expressed as a 150 kDa protein in various human tissues, including thymus, testis, and colon. SMC3 is expressed as a 146 kDa nuclear protein in the colon, but can also occur as a secreted proteoglycan (called Bamacan) expressed in testis and brain. Bamacan contains several glycosylation sites and is thought to play a role in basement membrane physiology.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 143 kDa

Swiss-Prot:

Q14683

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p-SMC1 (S957) polyclonal antibody detects endogenous levels of SMC1 protein only when phosphorylated at Ser957

DATA:




Western blot (WB) analysis of p-SMC 1 (S957) pAb at 1:500 dilution
Lane1:SGC7901 whole cell lysate(40ug)

Lane2:SGC7901 treated with UV for 5 minutes then repair for 1 hours
whole cell lysate(40ug)

Lane3:SGC7901 treated with UV for 5 minutes then repair for 3 hours
whole cell lysate(40ug)

Lane4:SGC7901 treated with UV for 5 minutes then repair for 16 hours
whole cell lysate(40ug)



Immunohistochemistry (IHC) analysis of p-SMC1 (S957) pAb in paraffin-embedded human lung carcinoma tissue.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology, Inc.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151