

## Stat3 (phospho-S727) polyclonal antibody

Catalog: BS4180

Host: Rabbit

Reactivity: Human, Mouse, Rat

### Background:

Membrane receptor signaling by various ligands, including interferons and growth hormones such as EGF, induces activation of JAK kinases which then leads to tyrosine phosphorylation of the various Stat transcription factors. Stat1 and Stat2 are induced by IFN- $\alpha$  and form a heterodimer which is part of the ISGF3 transcription factor complex. Although early reports indicate Stat3 activation by EGF and IL-6, it has been shown that Stat3 $\beta$  appears to be activated by both while Stat3 $\alpha$  is activated by EGF, but not by IL-6. Highest expression of Stat4 is seen in testis and myeloid cells. IL-12 has been identified as an activator of Stat4. Stat5 has been shown to be activated by Prolactin and by IL-3. Stat6 is involved in IL-4 activated signaling pathways.

### Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### Molecular Weight:

~ 88 kDa

### Swiss-Prot:

P40763

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IP: 1:50~1:200

### Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Specificity:

p-Stat3 (S727) polyclonal antibody detects endogenous levels of Stat3 protein only when phosphorylated at

Ser727.

### DATA:



Western blot (WB) analysis of

Stat3 (phospho-S727) polyclonal antibody at 1:500 dilution


Lane1:HepG2 whole cell lysate(40  $\mu$ g)

Lane2:A549 whole cell lysate(40  $\mu$ g)


Lane3:L02 whole cell lysate(40  $\mu$ g)

Lane4:PC12 whole cell lysate(40  $\mu$ g)

Lane5:3T3-L1 whole cell lysate(40  $\mu$ g)



Immunohistochemistry (IHC) analyzes of p-Stat3 (S727) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.



Western blot (WB) analysis of Stat3 (phospho-S727) pAb at 1:500 dilution

Lane1:HEK293T whole cell lysate

Lane2:HEK293T treated with TNF $\alpha$ (20ng/ml)for 5 minutes whole cell

### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151



## PRODUCT DATA SHEET

Bioworld Technology, Inc.

lysate

Lane3:HEK293T treated with TNF $\alpha$ (20ng/ml)for 20 minutes whole cell  
lysate

Lane4:HEK293T treated with TNF $\alpha$ (20ng/ml)for 40 minutes whole cell  
lysate

**Note:**

For research use only, not for use in diagnostic procedure.

---

**Bioworld Technology, Inc.**

**Add:** 1660 South Highway 100, Suite 500 St. Louis Park,  
MN 55416, USA.

**Email:** [info@bioworld.com](mailto:info@bioworld.com)

**Tel:** 6123263284

**Fax:** 6122933841

**Bioworld technology, co. Ltd.**

**Add:** No 9, weidi road Qixia District Nanjing, 210046,  
P. R. China.

**Email:** [info@biogot.com](mailto:info@biogot.com)

**Tel:** 0086-025-68037686

**Fax:** 0086-025-68035151