

CD26 Recombinant Rabbit mAb

Catalog: BS46021

Host: Rabbit

Reactivity: Mouse, Rat

BackGround:

Cell surface glycoprotein receptor involved in the costimulatory signal essential for T-cell receptor (TCR)-mediated T-cell activation. Acts as a positive regulator of T-cell coactivation, by binding at least ADA, CAV1, IGF2R, and PTPRC. Its binding to CAV1 and CARD11 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner. Its with interaction ADA also regulates lymphocyte-epithelial cell adhesion. In association with FAP is involved in the pericellular proteolysis of the extracellular matrix (ECM), the migration and invasion of endothelial cells into the ECM. May be involved in the promotion of lymphatic endothelial cells adhesion, migration and tube formation.

Product:

Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.

Molecular Weight:

120 kDa

Swiss-Prot:

P28843

Purification&Purity:

Affinity Purification

Applications:

WB: 1:1000
IHC: 1:200-1:1000


Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Isotype:

IgG

DATA:



Western blot analysis of extracts from Mouse thymus tissue at 1:1000.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 Fax: 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151