

## PP2A- $\alpha$ polyclonal antibody

Catalog: BS4867N

Host: Rabbit

Reactivity: Rat,Mouse,Human

### BackGround:

PP2A is the major phosphatase for microtubule-associated proteins (MAPs).PP2A can modulate the activity of phosphorylase B kinase casein kinase 2, mitogen-stimulated S6 kinase, and MAP-2 kinase.Cooperates with SGO2 to protect centromeric cohesin from separase-mediated cleavage in oocytes specifically during meiosis I (By similarity).Can dephosphorylate SV40 large T antigen and p53/TP53.Activates RAF1 by dephosphorylating it at 'Ser-259'.Mediates dephosphorylation of WEE1, preventing its ubiquitin-mediated proteolysis, increasing WEE1 protein levels, and promoting the G2/M checkpoint.Mediates dephosphorylation of MYC; promoting its ubiquitin-mediated proteolysis: interaction with AMBRA1 enhances interaction between PPP2CA and MYC.Mediates dephosphorylation of FOXO3; promoting its stabilization: interaction with AMBRA1 enhances interaction between PPP2CA and FOXO3.

### Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

### Molecular Weight:

~ 35 kDa

### Swiss-Prot:

P67775

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:1000~1:2000

IP: 1:50~1:200

### Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long

term. Avoid freeze-thaw cycles.

### Specificity:

PP2A- $\alpha$  polyclonal antibody detects endogenous levels of PP2A- $\alpha$  protein.

### DATA:




Western blot (WB) analysis of PP2A- $\alpha$  polyclonal antibody at 1:1000 dilution

Lane1:The liver tissue lysate of mouse(30ug)

Lane2:The heart tissue lysate of Rat(30ug)

Lane3:K562 whole cell lysate(30ug)



Immunoprecipitation of Mouse Liver lysates using PP2A- $\alpha$  pAb (Sepharose Bead Conjugate)#BD0048 (lane 2 and lane 3) and Nonspecific IgG Control (Sepharose Bead Conjugate)#BD0048 (lane 4) .Lane 1 is 30% input. The western blot was probed using PP2A- $\alpha$  pAb.

### Note:

For research use only, not for use in diagnostic procedure.

### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151