

HDAC4 Recombinant Rabbit mAb

Catalog:	BS49344	Host:	Rabbit	Reactivity:	Human, Mouse
BackGround:				Purification&Purity	:
Histones play a critical role in transcriptional regulation,				Affinity Purification	
cell cycle progression, and developmental events. Histone				Applications:	
acetylation/deacetylation alters chromosome structure and				WB: 1:1000	
affects transcription factor access to DNA. The protein				Storage&Stability:	
encoded by this gene belongs to class II of the histone				Store at 4 ${}^\circ\!\!{\rm C}$ short term. Aliquot and store at -20 ${}^\circ\!\!{\rm C}$ long	
deacetylase/acuc/apha family. It possesses histone				term. Avoid freeze-thaw cycles.	
deacetylase activity and represses transcription when			1	Isotype:	
tethered to a promoter. This protein does not bind DNA				IgG	
directly, but through transcription factors MEF2C and				DATA:	
MEF2D. It seems to interact in a multiprotein complex					
-	48 and HDAC3. [pro	vided by	RefSeq, Jul		HCT116
2008]				250	
Product:				1001	KD-
Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4),			.	70	<
0.15M NaCl, 40% Glycerol, 0.01% sodium azide and				50	<d td="" —<=""></d>
0.05% BSA. Stable for 12 months from date of receipt.			of receipt.		
Molecular Weight:				Western blot analysis of extracts from HCT116 cells using BS49344 at	
140 kDa				1: 1000.	
Swiss-Prot:				Note:	
P56524, Q6NZM9				For research use only, not for use in diagnostic procedure.	

Biowo	orld Technology, Inc.		
Add:	1660 South Highway 100, Suite 500 St. Louis Park,		
	MN 55416,USA.		
Email:	info@bioworlde.com		
Tel:	6123263284		
Fax:	6122933841		