

## **ARFGEF2** Recombinant Rabbit mAb

by this gene is involved in the activation of ARFs by accelerating replacement of bound GDP with GTP and is involved in Golgi transport. It contains a Sec7 domain, which may be responsible for its guanine-nucleotide exchange activity and also brefeldin A inhibition. [provided by RefSeq, Jul 2008] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight:	Stability:	Т24
in intracellular vesicular trafficking. The protein encoded by this gene is involved in the activation of ARFs by ac- celerating replacement of bound GDP with GTP and is involved in Golgi transport. It contains a Sec7 domain, which may be responsible for its guanine-nucleotide ex- change activity and also brefeldin A inhibition. [provided by RefSeq, Jul 2008] <b>Product:</b> Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. <b>Molecular Weight:</b>	℃ short terr d freeze-thav	w cycles.
by this gene is involved in the activation of ARFs by accelerating replacement of bound GDP with GTP and is involved in Golgi transport. It contains a Sec7 domain, which may be responsible for its guanine-nucleotide exchange activity and also brefeldin A inhibition. [provided by RefSeq, Jul 2008] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight:	d freeze-thav	w cycles.
Isotype:celerating replacement of bound GDP with GTP and is involved in Golgi transport. It contains a Sec7 domain, which may be responsible for its guanine-nucleotide ex- change activity and also brefeldin A inhibition. [provided by RefSeq, Jul 2008]Isotype: IgGProduct: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight:Isotype: 		Т24
involved in Golgi transport. It contains a Sec7 domain, which may be responsible for its guanine-nucleotide ex- change activity and also brefeldin A inhibition. [provided by RefSeq, Jul 2008] <b>Product:</b> Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. <b>Molecular Weight:</b>	кDа	
which may be responsible for its guanine-nucleotide exchange activity and also brefeldin A inhibition. [provided by RefSeq, Jul 2008] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight:	kDa	
change activity and also brefeldin A inhibition. [provided by RefSeq, Jul 2008] <b>Product:</b> Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. <b>Molecular Weight:</b>	kDa	
by RefSeq, Jul 2008] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight:	kDa	
Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight:	<b>kD</b> a	a
Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight:	250	
0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight:	100	-
0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight:	70 50 40	E
Molecular Weight:	35	=
	18	=
	analysis of extra	racts from T24 cells using BS49483 at 1:
Swiss-Prot: 3000.		
Q9Y6D5 Note:		
Purification&Purity: For researce	ch use only, i	not for use in diagnostic procedure.
Affinity Purification		
Applications:		

## Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151