

Brn-3c polyclonal antibody

Cuturog	Catal	log:	BS5632
---------	-------	------	--------

Host: Ra

Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Three Brn-3 proteins have been described and are designated Brn-3a, Brn-3b and Brn-3c. Brn-3a has two functional transactivating domains, one at the amino terminus and one at the carboxy terminus. While Brn-3a and Brn-3c stimulate transcription, Brn-3b generally functions as a transcriptional repressor. However, Brn-3b, but not Brn-3a, has been shown to regulate the expression of the acetylcholine receptor.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 37 kDa

Swiss-Prot:

Q15319

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Specificity:

Brn-3c polyclonal antibody detects endogenous levels of Brn-3c protein.

DATA:



Western blot (WB) analysis of Brn-3c polyclonal antibody at 1:500 dilu-

tion

Lane1:Hela cell lysate

Lane2:Raw264.7 cell lysate

Lane3:H9C2 cell lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151