

PIK3C2G polyclonal antibody

Catalog: BS60313

Host: Rabbit

Reactivity: Human

BackGround:

Phosphoinositide 3-kinase activity is implicated in assorted cellular responses activated by mammalian cell surface receptors and the regulation of protein sorting in yeast. The p110 γ (PIK3CG) enzyme is activated in vitro by both the α and $\beta\gamma$ subunits of heterotrimeric GTP-binding proteins (G proteins) and does not associate with a p85 adaptor molecule. PI 3-kinase C2 γ , also designated p110 γ , may link signaling through G protein-coupled receptors to the generation of phosphoinositide second messengers that are phosphorylated in the D-3 position. The PI 3-kinase C2 γ gene encodes a 1,050 amino acid polypeptide with 36% identity to human PI 3-kinase C2 α . Research indicates that PI 3-kinase C2 γ can block the growth of human colon cancer cells.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 171 kDa

Swiss-Prot:

O75747

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

PIK3C2G polyclonal antibody detects endogenous levels of PIK3C2G protein.

DATA:



Western blot (WB) analysis of PIK3C2G polyclonal antibody at 1:500 dilution

Lane1:PC3 whole cell lysate(40 μ g)

Lane2:HepG2 whole cell lysate(40 μ g)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151