

## TREM1 polyclonal antibody

Catalog: BS60555-B

Host: Rabbit

Reactivity: Human

### BackGround:

TREM-1 (triggering receptor expressed on myeloid cells-1) is expressed in monocytes and neutrophils but not in lymphocytes, dendritic cells, or other cell types. TREM-1 is a glycoprotein that is reduced by deglycosylation, in agreement with the predicted molecular mass. TREM-1 is an activating receptor of the Ig superfamily that is expressed on human myeloid cells, selectively expressed on blood neutrophils and a subset of monocytes, and is upregulated by bacterial LPS. Immunoblot analysis shows that TREM-1 is associated with DAP12, a molecule frequently associated with activating receptors. TREM-1 and the myeloid DAP12-associating lectin (MDL-1) are recently identified receptors which associate non-covalently with DAP12 to form receptor complexes that are involved in monocytic activation and inflammatory response.

### Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

### Molecular Weight:

~ 25kDa

### Swiss-Prot:

Q9NP99

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:2000~1:5000


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Specificity:

TREM1 polyclonal antibody detects endogenous levels of TREM1 protein.

### DATA:



Western blot (WB) analysis of TREM1 polyclonal antibody at 1:2000 dilution

Lane1:The spleen tissue lysate of Rat(40ug)

Lane2:The spleen tissue lysate of mouse(40ug)


Lane3:HepG2 whole cell lysate(40ug)

Lane4:A549 whole cell lysate(40ug)

Lane5:THP-1 whole cell lysate(40ug)

Lane6:PC12 whole cell lysate(40ug)

Lane7:Raw264.7 whole cell lysate(40ug)



Immunoprecipitation of HepG2 cell lysates using TREM1 pAb (Sepharose Bead Conjugate)#BD0048 (lane 3 and lane 4) and Nonspecific IgG Control (Sepharose Bead Conjugate)#BD0048 (lane 1 and lane 2). Lane 5 is 30% input. The western blot was probed using TREM1 pAb.

### Note:

For research use only, not for use in diagnostic procedure.

### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151



## PRODUCT DATA SHEET

Bioworld Biotech Co., Ltd

---

---

### **Bioworld Technology, Inc.**

Add: 1660 South Highway 100, Suite 500 St. Louis Park,  
MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

### **Bioworld technology, co. Ltd.**

Add: No 9, weidi road Qixia District Nanjing, 210046,  
P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151