

ATP6V0A2 polyclonal antibody

Catalog: **BS60737** Host:

Rabbit

Reactivity: Human, Mouse

BackGround:

Vacuolar-type H+-ATPase (V-ATPase) is a multisubunit enzyme responsible for the acidification of eukaryotic intracellular organelles. V-ATPases pump protons against an electrochemical gradient, while F-ATPases reverse the process, thereby synthesizing ATP. A peripheral V1 domain, which is responsible for ATP hydrolysis, and an integral V0 domain, which is responsible for proton translocation, comprise the V-ATPase complex. Nine subunits (A-H) make up the V1 domain and five subunits (A, D, C, C' and C") make up the V0 domain. As part of the V0 domain, V-ATPase A2 (ATPase, H+ transporting, lysosomal V0 subunit a2), consists of 856 amino acids and is also known as ATP6V0A2, V-type proton ATPase subunit a isoform 2, vacuolar proton translocating ATPase subunit a isoform 2, lysosomal H(+)-transporting ATPase V0 subunit a2 or TJ6. V-ATPase A2 is a multi-pass membrane protein with localization in the cell membrane, endosome membrane and the subapical vesicles of the kidney's proximal tubules. V-ATPase A2 plays an important role in Golgi function by regulating pH. Wrinkly skin syndrome (WSS) and cutis laxa type II (ARCL type II) are caused as a result of V-ATPase A2 defects.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

- **Molecular Weight:**
- ~ 98 kDa

Swiss-Prot:

O9Y487

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 102% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:207


Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-27 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Specificity:

ATP6V0A2 polyclonal antibody detects endogenous levels of ATP6V0A2 protein.

DATA:



Western blot (WB) analysis of ATP6V0A2 polyclonal antibody at 1:500 dilution

Lane1:HEK293T whole cell lysate

Lane2:sp2/0 whole cell lysate

Lane3:H9C9 whole cell lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc. Add:

1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. **Email:** info@bioworlde.com Tel: 6123263284 6122933841 Fax:

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. **Email:** info@biogot.com Tel: 0086-025-68037686 0086-025-68035151 Fax: