

NFAT3 (Phospho-S676) polyclonal antibody

Catalog: BS64503

Host: Rabbit

Reactivity: Human

BackGround:

NFATc4 (nuclear factor of activated T cells, cytoplasmic, calcineurin-dependent 4) is a member of the nuclear factors of activated T cells DNA-binding transcription complex that influences cytokine gene expression, cardiac hypertrophy and adipocyte differentiation. This complex consists of at least two components: a cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation and an inducible nuclear component. Other members of this family participate in the formation of this complex. NFATc4 plays a role in the inducible expression of cytokine genes in T cells, including the induction of IL-2 and IL-4. p38 MAP kinase phosphorylates multiple residues in the NFAT homology domain of NFATc4.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 140 kDa

Swiss-Prot:

Q14934

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB:1:500~1:1000 IHC:1:50~1:200 IF:1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

NFAT3 (Phospho-S676) polyclonal antibody detects endogenous levels of NFAT3 protein only when phosphorylated at Ser676.

DATA:



Western blot (WB) analysis of NFAT3 (Phospho-S676) polyclonal antibody at 1:500 dilution

LaneA:The Testis tissue lysate of Mouse

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151