

DTX3L Polyclonal Antibody

Catalog: BS65777

Host: Rabbit

Reactivity: Human, Mouse, Rat, Rabbit,

BackGround:

DTX3L functions as an E3 ubiquitin ligase

Product:

0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

Molecular Weight:

83kDa

Swiss-Prot:

Q8TDB6

Purification&Purity:

affinity purified by Protein A

Applications:

IHC-P=1:100-500


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:


DTX3L Polyclonal Antibody detects endogenous levels of DTX3L protein.

DATA:



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block

endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 °C for 30min; Antibody incubation with (DTX3L) Polyclonal Antibody, Unconjugated (bs-14438R) at 1:200 overnight at 4 °C



Tissue/cell: human kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37 °C for 20 min; Incubation: Anti-DTX3L Polyclonal Antibody, Unconjugated 1:200, overnight at 4 °C,

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151