

Tartrate Resistant Acid Phosphatase Polyclonal Antibody

Catalog: BS65953	Host: Rabbit	Reactivity: Hu-man, Mouse, Rat, Dog, Pig, Horse, Rabbit,
------------------	--------------	--

BackGround:

Tartrate-resistant acid phosphatase is an iron containing glycoprotein that catalyzes the conversion of orthophosphoric monoester to alcohol and orthophosphate. TRAP is the most basic of the acid phosphatases and is the only form not inhibited by L+-tartrate. TRAP is a relatively minor lysosomal enzyme which may be activated in certain pathologies such as Hodgkin's disease and B- and T-cell leukemias. Receptor activator of NFκB ligand (RANKL) plays an essential role in osteoclast differentiation and activation by increasing the expression of protease osteoclast markers such as TRAP. TRAP has collagenolytic activity and plays a major role in ligament degradation.

Product:

0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

Molecular Weight:

35 kD

Swiss-Prot:

P13686

Purification&Purity:

affinity purified by Protein A

Applications:

WB=1:500-2000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.


Specificity:

Tartrate Resistant Acid Phosphatase Polyclonal Antibody detects endogenous levels of Tartrate Resistant Acid Phosphatase protein.

DATA:



Primary: Anti-Tartrate Resistant Acid Phosphatase at 1/1000 dilution



Primary: Anti-Tartrate Resistant Acid Phosphatase at 1/300 dilution

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151