

Monoclonal Anti-MMAE specific Antibody, Rabbit IgG (M1D12)

Catalog: BS66200

Host: Rabbit

Reactivity:

BackGround:

Monomethyl auristatin E (MMAE) is a synthetic derivative of dolastatin 10 and functions as a potent mitotic inhibitor by inhibiting tubulin polymerization. MMAE is widely used as a cytotoxic component of antibody-drug conjugates (ADCs) to treat several cancer types. Anti-MMAE antibody is a rabbit monoclonal antibody specially react with MMAE without MMAF, which is more sensitive than mouse antibody. The anti-MMAE antibody is a useful reagent in PK assay to determine conjugated antibodies.

Product:

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Molecular Weight:

Swiss-Prot:

Purification&Purity:

Protein A purified / Protein G purified

Applications:

ELISA 0.06-125 ng/mL


Storage&Stability:

-20 $^{\circ}$ C to -70 $^{\circ}$ C for 12 months in lyophilized state;
-70 $^{\circ}$ C for 3 months under sterile conditions after reconstitution.


Specificity:

Specifically recognizes MMAE and does not recognize MMAF.


DATA:



Immobilized Disitamab Vedotin (RC48) at 0.2 μ g/mL (100 μ L/well) can bind Monoclonal Anti-MMAE specific Antibody, Rabbit IgG (M1D12) with a linear range of 0.06-2 ng/mL (QC tested).



Immobilized Monoclonal Anti-MMAE specific Antibody, Rabbit IgG (M1D12) at 5 μ g/mL, add Disitamab Vedotin (RC48) in the 100% Human Serum and then add Biotinylated Human Her2, His,Avitag, premium grade at 0.5 μ g/mL. Detection was performed using HRP-conjugated Streptavidin (QC tested).



Immobilized Monoclonal Anti-MMAE specific Antibody, Rabbit IgG (M1D12) at 5 μ g/mL, add Disitamab Vedotin (RC48) and then add Biotinylated Human Her2, His,Avitag, premium grade at 0.5 μ g/mL. Detection was performed using HRP-conjugated Streptavidin (Routinely tested).

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.


Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



Inhibition of Monoclonal Anti-MMAE specific Antibody, Rabbit IgG (M1D12) ELISA
0.1 µg of Monoclonal Anti-MMAE specific Antibody, Rabbit IgG (M1D12) per well



Note:

For research use only, not for use in diagnostic procedure.

Serial dilutions of Monoclonal Anti-MMAE specific Antibody, Rabbit IgG (M1D12): Disitamab Vedotin (RC48) binding reactions. The half maximal inhibitory concentration (IC50) is 0.006187 µg/mL (Routinely tested).

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151