

Histone H3 (TriMethyl-K9) polyclonal antibody

Catalog:	BS67150	Host:	Rabbit	Reactivity:	Human,Rat,Mouse
BackGround:				WB (1/500 - 1/1000)	
Core component of nucleosome. Nucleosomes wrap and				Storage&Stability:	
compact DNA into chromatin, limiting DNA accessibility				Store at 4 ${}^\circ\!\!{\rm C}$ short term. Aliquot and store at -20 ${}^\circ\!\!{\rm C}$ long	
to the cellular machineries which require DNA as a tem-				term. Avoid freeze-thaw cycles.	
plate. Histones thereby play a central role in transcription				Specificity:	
regulation, DNA repair, DNA replication and chromoso-				Recognizes endogenous levels of Histone H3 with a site	
mal stability. DNA accessibility is regulated via a com-				at TriMethyl K9 protein.	
plex set of post-translational modifications of histones,			of histones,	DATA:	
also called histone code, and nucleosome remodeling.			odeling.	kDa A B C	
Product:				34	
Liquid in 0.42% Potassium phosphate, 0.87% Sodium			7% Sodium	26	
chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.			dium azide.		
Molecular Weight:				17	
~ 17 kDa				10	
Swiss-Prot:	;				1
P68431; Q71DI3; P84243; Q6NXT2				Western blot analysis of Histone H3 (TriMethyl K9) expression in	
Purification&Purity:				HEK293T (A), A549 (B), H1688 (C) whole cell lysates.	
The antibody was affinity-purified from rabbit antiserum			it antiserum	Note:	
by affinity-chromatography using epitope-specific im-			specific im-	For research use only, not for use in diagnostic procedure.	

Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 Fax: 6122933841

munogen and the purity is > 111% (by SDS-PAGE).

Applications:

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151