

BS67157

Host:

Rabbit

Catalog:

Applications:

Human,Rat,Mouse

Histone H3 (MonoMethyl-K122) polyclonal antibody

Reactivity:

BackGround:	WB (1/500 - 1/1000)
Core component of nucleosome. Nucleosomes wrap and	Storage&Stability:
compact DNA into chromatin, limiting DNA accessibility	Store at 4 ${}^\circ\!\!{\rm C}$ short term. Aliquot and store at -20 ${}^\circ\!\!{\rm C}$ long
to the cellular machineries which require DNA as a tem-	term. Avoid freeze-thaw cycles.
plate. Histones thereby play a central role in transcription	Specificity:
regulation, DNA repair, DNA replication and chromoso-	Recognizes endogenous levels of Histone H3 with a site
mal stability. DNA accessibility is regulated via a com-	at MonoMethyl-K122 protein.
plex set of post-translational modifications of histones,	DATA:
also called histone code, and nucleosome remodeling.	kDa A B
Product:	34
Liquid in 0.42% Potassium phosphate, 0.87% Sodium	26
chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.	
Molecular Weight:	17
~ 17 kDa	10
Swiss-Prot:	
P68431; Q71DI3; P84243; Q6NXT2	Western blot analysis of Histone H3 (MonoMethyl-K122) expression in
Purification&Purity:	mouse testis (A), mouse muscle (B) whole cell lysates.
The antibody was affinity-purified from rabbit antiserum	Note:
by affinity-chromatography using epitope-specific im-	For research use only, not for use in diagnostic procedure.
munogen and the purity is $> 118\%$ (by SDS-PAGE).	

Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 Fax: 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. **Email:** info@biogot.com 0086-025-68037686 Tel: 0086-025-68035151 Fax: