

BS67159

Host:

Rabbit

Catalog:

Human,Rat,Mouse

Histone H4 (MonoMethyl-K12) polyclonal antibody

Reactivity:

<u> </u>	·
BackGround:	WB (1/500 - 1/1000)
Core component of nucleosome. Nucleosomes wrap and	Storage&Stability:
compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a tem-	Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.
plate. Histones thereby play a central role in transcription	Specificity:
regulation, DNA repair, DNA replication and chromoso- mal stability. DNA accessibility is regulated via a com-	Recognizes endogenous levels of Histone H4 with a site at MonoMethyl-K12 protein.
plex set of post-translational modifications of histones,	DATA:
also called histone code, and nucleosome remodeling.	kDa A B C
Product:	26
Liquid in 0.42% Potassium phosphate, 0.87% Sodium	20
chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide. Molecular Weight:	17
~ 17 kDa	10
Swiss-Prot:	
P62805	Western blot analysis of Histone H4 (MonoMethyl-K12) expression in
Purification&Purity:	HEK293T (A), H446 (B), H1688 (C) whole cell lysates.
The antibody was affinity-purified from rabbit antiserum	Note:

For research use only, not for use in diagnostic procedure.

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 120% (by SDS-PAGE).

Applications:

Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 Fax: 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151