

**Bioworld Technology,Inc.** 

# **SIRT3 polyclonal antibody**

Catalog: BS67761

Host: F

Rabbit

Reactivity: Human, Mouse, Rat

# **BackGround:**

The Silent Information Regulator (SIR2) family of genes is a highly conserved group of genes that encode nicotinamide adenine dinucleotide (NAD)-dependent protein deacetylases, also known as Class III histone deacetylases. The first discovered and best characterized of these genes is Saccharomyces cerevisiae Sir2, which is involved in silencing of mating type loci, telomere maintenance, DNA damage response and cell aging. SirT3, a mammalian homolog of Sir2, exists in human cells in two forms. The full-length 44 kDa protein localizes to the nucleus, while a processed 28 kDa protein lacking 142 amino terminal residues localizes exclusively to the mitochondria. The single murine form of SirT3 is equivalent to the processed human SirT3 protein. Full-length SirT3 protein is processed in the mitochondrial matrix by the mitochondrial matrix processing peptidase (MMP). Both full-length and processed forms of SirT3 are enzymatically active and de-acetylate histone H3 at Lys9 and histone H4 at Lys16 in vitro. SirT3 also de-acetylates Lys642 of acetyl-CoA synthetase 2 (AceCS2) and activates AceCS2 activity in the mitochondria. Restricted caloric intake, which is linked to increased lifespan in multiple organisms, increases SirT3 expression in white and brown adipocytes of obese mice, suggesting a role for SirT3 in aging. Two observations implicate SirT3 in the regulation of mitochondrial thermogenesis. First, exposure to cold temperatures increases SirT3 expression in brown adipocytes, while elevated temperatures reduce SirT3 expression. Second, over-expression of SirT3 results in increased levels of the mitochondrial uncoupling protein 1 (UCP1). SirT3 protein levels are also elevated in certain breast cancers.

#### **Product:**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Bioworld Technology, Inc.			
Add:	1660 South Highway 100, Suite 500 St. Louis Park,		
	MN 55416,USA.		
Email:	info@bioworlde.com		
Tel:	6123263284		
Fax:	6122933841		

#### **Molecular Weight:**

~ 29 kDa

**Swiss-Prot:** 

O9NTG7

#### **Purification&Purity:**

The antibody was purified by immunogen affinity chromatography.

## **Applications:**

WB (1/500 - 1/1000), IHC (1/50 - 1/200), IF/ICC (1/50 - 1/200)

#### **Storage&Stability:**

K

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.

## **Specificity:**

Recognizes endogenous levels of SIRT3 protein.

## DATA:

Da	А	в	С
95			
72			
55			
43			
34			
26		-	-
17			

Western blot analysis of SIRT3 expression in HepG2 (A), mouse liver (B), rat liver (C) whole cell lysates.



Immunohistochemical analysis of SIRT3 staining in rat kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature

Bioworld technology, co. Ltd.Add:No 9, weidi road Qixia District Nanjing, 210046,<br/>P. R. China.Email:info@biogot.comTel:0086-025-68037686Fax:0086-025-68035151



# **PRODUCT DATA SHEET**

## **Bioworld Technology,Inc.**

and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of SIRT3 staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Note:

For research use only, not for use in diagnostic procedure.

DIOWO	niu rechnology, nic.
Add:	1660 South Highway 100, Suite 500 St. Louis Park
	MN 55416,USA.
Email:	info@bioworlde.com

Disward Tashnalagu Ing

 $\begin{array}{c} 6123263284\\ 6122933841 \end{array}$ 

Tel:

Fax:

# Bioworld technology, co. Ltd.

Add:	No 9, weidi road Qixia District Nanjing, 210046
	P. R. China.
Email:	info@biogot.com
Tel:	0086-025-68037686
Fax:	0086-025-68035151