

# [KO Validated] NUCB1 polyclonal antibody

Catalog: BS71997

Host: Ral

```
Rabbit
```


Human, Mouse, Rat

ECL Basic Kit .< br/>Exposure time: 90s.


Reactivity:

# Control Contr

Western blot analysis of extracts from normal and NUCB1 knockout 293T cells, using NUCB1 antibody at 1:3000 dilution.<br/>br/>Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>br/>Lysates/proteins: 25ug per lane.<br/>br/>Blocking buffer: 3% nonfat dry milk in TBST.<br/>br/>Detection: ECL Basic Kit .<br/>br/>Exposure time: 3s.



Immunofluorescence analysis of L929 cells using NUCB1 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using NUCB1 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

#### Note:

For research use only, not for use in diagnostic procedure.

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151

# **BackGround:**

This gene encodes a member of a small calcium-binding EF-hand protein family. The encoded protein is thought to have a key role in Golgi calcium homeostasis and Ca(2+)-regulated signal transduction events.

### **Product:**

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

**Molecular Weight:** 

60kDa

**Swiss-Prot:** 

Q02818

**Purification&Purity:** 

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

# **Applications:**

WB,1:500 - 1:2000|IHC,1:50 - 1:100|IF/ICC,1:50 - 1:100


#### Storage&Stability:

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at -20  $^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.

#### **Category:**

**Polyclonal Antibodies** 

## **DATA:**



Western blot analysis of extracts of various cell lines, using NUCB1 antibody at 1:3000 dilution.<br/>br/>Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>br/>Lysates/proteins: 25ug per lane.<br/>br/>Blocking buffer: 3% nonfat dry milk in TBST.<br/>betection:

# Bioworld Technology, Inc.

 
 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841