

## **Rock-2 polyclonal antibody**

| Catalog: BS7216 Host: Rabb   | it Reactivity: Human, Mouse, Rat  |
|--|---|
| BackGround:<br>Rho proteins function as molecular switches that are<br>to turn cytokinesis on and off. Although little is ka<br>about signaling downstream of Rho, a host of put<br>Rho effector proteins have been de-scribed, inclu-<br>rhophilin, Rhotekin, citron and the serine/threonine<br>nase, protein kinase N. Two additional Rho-activated<br>ine/threonine kinases have been described, desig<br>Rock-1 and Rock-2 (also referred to as Roka<br>Rho-associated coil-containing protein kinase. Re<br>and Rock-2 share a structural similarity with myo<br>dystrophy kinase.<br>Product: | hown by affinity-chromatography using epitope-specific im-<br>munogen and the purity is > 95% (by SDS-PAGE).<br>Applications:<br>e ki-<br>WB: 1:500~1:2000<br>d ser-<br>Storage&Stability:<br>nated Store at 4 $^{\circ}$ short term. Aliquot and store at -20 $^{\circ}$ long<br>term. Avoid freeze-thaw cycles.<br>Specificity:<br>tonic Rock-2 polyclonal antibody detects endogenous levels of<br>Rock-2 protein. |
| Rabbit IgG, 1mg/ml in PBS with 0.02% sodium a  | DATA:<br>uzide,   |
| 50% glycerol, pH7.2<br>Molecular Weight:<br>~ 160 kDa<br>Swiss-Prot:<br>O75116   | Western blot (WB) analysis of Rock-2 pAb at 1:500 dilution<br>LaneA:The Liver tissue lysate of Mouse(40ug)<br>LaneB:The Liver tissue lysate of Rat(40ug)<br>Note:<br>For research use only, not for use in diagnostic procedure.  |

## Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: <u>info@bioworlde.com</u> Tel: 6123263284 Fax: 6122933841