

# Phospho-PKC (pan) (zeta)-T410 polyclonal antibody

Catalog: BS74146

Host: Rabbit

Reactivity: Human, Mouse, Rat

### **BackGround:**

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This kinase has been reported to play roles in many different cellular processes, such as cell adhesion, cell transformation, cell cycle checkpoint, and cell volume control. Knockout studies in mice suggest that this kinase may be a fundamental regulator of cardiac contractility and Ca(2+) handling in myocytes. [provided by RefSeq, Jul 2008]

#### **Product:**

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

#### **Molecular Weight:**

#### 85kDa

**Swiss-Prot:** 

P17252/P05771/Q05655/Q02156/P05129 /P24723/P41743/Q04759/Q05513

#### **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

**Applications:** 

WB,1:500 - 1:2000


**Storage&Stability:** 

Store at  $4 \,^{\circ}$  short term. Aliquot and store at  $-20 \,^{\circ}$  long term. Avoid freeze-thaw cycles.

**Modification:** 

Phosphorylated

#### **DATA:**



Western blot analysis of extracts of C6 cells, using Phospho-PKC -T410 antibody at 1:1000 dilution. C6 cells were treated by PMA/TPA at 37°C for 30 minutes after serum-starvation overnight.<br/>br/>Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>br/>Lysates/proteins: 25ug per lane.<br/>br/>Blocking buffer: 3% BSA.<br/>br/>Detection: ECL Basic Kit .<br/>br/>Exposure time: 90s.

#### Note:

For research use only, not for use in diagnostic procedure.

#### Bioworld Technology, Inc.

 
 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

## Bioworld technology, co. Ltd.

 
 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151