

## PRODUCT DATA SHEET

Bioworld Technology,Inc.

# Phospho-SHP1-Y564 polyclonal antibody

Catalog: BS74171 Host: Rabbit Reactivity: Human, Mouse

#### **BackGround:**

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. N-terminal part of this PTP contains two tandem Src homolog (SH2) domains, which act as protein phospho-tyrosine binding domains, and mediate the interaction of this PTP with its substrates. This PTP is expressed primarily in hematopoietic cells, and functions as an important regulator of multiple signaling pathways in hematopoietic cells. This PTP has been shown to interact with, and dephosphorylate a wide spectrum of phospho-proteins involved in hematopoietic cell signaling. Multiple alternatively spliced variants of this gene, which encode distinct isoforms, have been reported.

## **Product:**

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

#### **Molecular Weight:**

60kDa

#### **Swiss-Prot:**

P29350

## **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

## **Applications:**

WB,1:500 - 1:2000


## Storage&Stability:

Store at  $4 \, \mathbb{C}$  short term. Aliquot and store at  $-20 \, \mathbb{C}$  long term. Avoid freeze-thaw cycles.

## **Modification:**

Phosphorylated

#### **DATA:**



Western blot analysis of extracts of HeLa and RAW 264.7 cells, using Phospho-SHP1-Y564 antibody at 1:2000 dilution. HeLa cell lysates were treated by CIP at 37 °C for 1 hour.RAW 264.7 cell lysates were treated by CIP at 37 °C for 1 hour.<br/>
Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
Springly-Sproteins: 25ug per lane.<br/>
Springly-Sproteins: 25ug per lane.<br/>
Springly-Sproteins: 25ug per lane.<br/>
Springly-Sproteins: 25ug per lane.<br/>
Springly-Sproteins: 3% BSA.<br/>
Springly-Sproteins: 25ug per lane.

#### Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u>
Tel: 0086-025-68037686
Fax: 0086-025-68035151