

[KO Validated] BRG1/SMARCA4 polyclonal antibody

Catalog: BS74306

Host: R

Rabbit

Reactivity: Human

BackGround:

The protein encoded by this gene is a member of the SWI/SNF family of proteins and is similar to the brahma protein of Drosophila. Members of this family have helicase and ATPase activities and are thought to regulate transcription of certain genes by altering the chromatin structure around those genes. The encoded protein is part of the large ATP-dependent chromatin remodeling complex SNF/SWI, which is required for transcriptional activation of genes normally repressed by chromatin. In addition, this protein can bind BRCA1, as well as regulate the expression of the tumorigenic protein CD44. Mutations in this gene cause rhabdoid tumor predisposition syndrome type 2. Multiple transcript variants encoding different isoforms have been found for this gene.

Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

180kDa

Swiss-Prot:

P51532

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB,1:500 - 1:2000|IF/ICC,1:20 - 1:50

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Modification:

Unmodification

DATA:


Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.


 Email:
 info@bioworlde.com

 Tel:
 6123263284


 Fax:
 6122933841



Western blot analysis of extracts of various cell lines, using BRG1/BRG1/SMARCA4 antibody at 1:500 dilution.
br/>Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.
br/>Lysates/proteins: 25ug per lane.
br/>Blocking buffer: 3% nonfat dry milk in TBST.



Western blot analysis of extracts from normal and BRG1/BRG1/SMARCA4 knockout 293T cells, using BRG1/BRG1/SMARCA4 antibody at 1:500 dilution.
br/>Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.
br/>Lysates/proteins: 25ug per lane.
br/>Blocking buffer: 3% nonfat dry milk in TBST.
br/>Detection: ECL Basic Kit .
br/>Exposure time: 90s.



Immunofluorescence analysis of HeLa cells using

BRG1/BRG1/SMARCA4 antibody .

Note:

Bioworld technology, co. Ltd.Add:No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.Email:info@biogot.comTel:0086-025-68037686Fax:0086-025-68035151



Bioworld Technology, Inc.

For research use only, not for use in diagnostic procedure.

Bioworld	Technol	loqy,	Inc.
----------	---------	-------	------

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 Fax: 6122933841

Bioworld technology, co. Ltd.

 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151