

polyclonal antibody HRH2

Catal	log:	BS75626
-------	------	---------

Host:

Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Histamine is a ubiquitous messenger molecule released from mast cells, enterochromaffin-like cells, and neurons. Its various actions are mediated by histamine receptors H1, H2, H3 and H4. Histamine receptor H2 belongs to the family 1 of G protein-coupled receptors. It is an integral membrane protein and stimulates gastric acid secretion. It also regulates gastrointestinal motility and intestinal secretion and is thought to be involved in regulating cell growth and differentiation. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

38kDa

Swiss-Prot:

P25021

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB,1:500 - 1:2000


Storage&Stability:

Store at $4 \, \mathbb{C}$ short term. Aliquot and store at $-20 \, \mathbb{C}$ long term. Avoid freeze-thaw cycles.

Modification:

Unmodification

DATA:



Western blot analysis of extracts of various cell lines, using HRH2 antibody at 1:1000 dilution.
Secondary antibody: HRP Goat Anat 1:10000 dilution.
br/>Lysates/proteins: 25ug per ti-Rabbit IgG lane.
br/>Blocking buffer: 3% nonfat dry milk in TBST.
br/>Detection: ECL Basic Kit .< br/>
Exposure time: 90s.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. **Email:** info@bioworlde.com Tel: 6123263284 6122933841 Fax:

Bioworld technology, co. Ltd. No 9, weidi road Qixia District Nanjing, 210046, Add: P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 0086-025-68035151 Fax: