

# DDX17 polyclonal antibody

Catalog: BS77079

Host: Ra

Rabbit

Reactivity: Human, Mouse, Rat

## **BackGround:**

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure, such as translation initiation, nuclear and mitochondrial splicing, and ribosome and splicesosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein, which is an ATPase activated by a variety of RNA species, but not by dsDNA. This protein, and that encoded by DDX5 gene, are more closely related to each other than to any other member of the DEAD box family. This gene can encode multiple isoforms due to both alternative splicing and the use of alternative translation initiation codons, including a non-AUG (CUG) start codon.

## **Product:**

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

## **Molecular Weight:**

80kDa

**Swiss-Prot:** 

#### Q92841

#### **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

## **Applications:**

WB,1:500 - 1:2000|IHC,1:50 - 1:200|IF/ICC,1:50 - 1:200


Storage&Stability:

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.


Modification:

Unmodification

**DATA:** 



Western blot analysis of extracts of various cell lines, using DDX17 antibody at 1:1000 dilution.<br/>br/>Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>br/>Lysates/proteins: 25ug per lane.<br/>br/>Blocking buffer: 3% nonfat dry milk in TBST.<br/>br/>Detection: ECL Basic Kit .<br/>br/>Exposure time: 5min.



Immunofluorescence analysis of U2OS cells using DDX17 antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

#### Note:

For research use only, not for use in diagnostic procedure.

## Bioworld Technology, Inc.

Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841

## Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151