

PRODUCT DATA SHEET

Bioworld Technology,Inc.

RPL35 polyclonal antibody

Catalog: BS77369 Host: Rabbit Reactivity: Human, Mouse, Rat

BackGround:

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L29P family of ribosomal proteins. It is located in the cytoplasm. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq, Jul 2008]

Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

30kDa

Swiss-Prot:

P42766

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB,1:500 - 1:2000|IHC,1:50 - 1:200|IF/ICC,1:50 - 1:200


Storage&Stability:

Store at $4\,\mathrm{C}$ short term. Aliquot and store at -20 C long term. Avoid freeze-thaw cycles.

Modification:

Unmodification

DATA:



Western blot analysis of extracts of various cell lines, using RPL35
Rabbit pAb at 1:1000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.


Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.


Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat
Anti-Rabbit IgG at 1:10000 dilution.



Immunohistochemistry of paraffin-embedded Human colon carcinoma using RPL35 Rabbit pAb at dilution of 1:100 .Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of L929 cells using RPL35 antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u>
Tel: 0086-025-68037686
Fax: 0086-025-68035151