

IL13RA2 polyclonal antibody

ti-Rabbit IgG

ECL Basic Kit .< br/>Exposure time: 180s.

Catalog: BS77895

Host: R

Rabbit

Reactivity: Human, Mouse, Rat

lane.
br/>Blocking buffer: 3% nonfat dry milk in TBST.
br/>Detection:

at 1:10000 dilution.
br/>Lysates/proteins: 25ug per

BackGround:

The protein encoded by this gene is closely related to II13RA1, a subuint of the interleukin 13 receptor complex. This protein binds IL13 with high affinity, but lacks cytoplasmic domain, and does not appear to function as a signal mediator. It is reported to play a role in the internalization of IL13.

Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

44kDa

Swiss-Prot:

Q14627

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB,1:500 - 1:2000|IHC,1:50 - 1:200|IF/ICC,1:50 - 1:200


Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Modification:

Unmodification

DATA:



Western blot analysis of extracts of various cell lines, using IL13RA2 antibody at 1:1000 dilution.
Secondary antibody: HRP Goat An-

Bioworld Technology, Inc.


 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com


 Tel:
 6123263284

 Fax:
 6122933841
 130kDa 100kDa 70kDa 55kDa 40kDa 25kDa 25kDa 25kDa

Western blot analysis of extracts of Rat liver, using IL13RA2 antibody at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.
br/>Lysates/proteins: 25ug per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
br/>Detection: ECL Basic Kit .
Exposure time: 90s.



Immunofluorescence analysis of C6 cells using IL13RA2 antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using IL13RA2 anti-

body at dilution of 1:100. Blue: DAPI for nuclear staining.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld technology, co. Ltd.
Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.
Email: info@biogot.com Tel: 0086-025-68037686
Fax: 0086-025-68035151