

OTULIN polyclonal antibody

Catalog: BS78035

Host: Rabbit

Reactivity: Hun

ity: Human, Mouse, Rat

BackGround:

This gene encodes a member of the peptidase C65 family of ubiquitin isopeptidases. Members of this family remove ubiquitin from proteins. The encoded enzyme specifically recognizes and removes M1(Met1)-linked, or linear, ubiquitin chains from protein substrates. Linear ubiquitin chains are known to regulate the NF-kappa B signaling pathway in the context of immunity and inflammation. Mutations in this gene cause a potentially fatal autoinflammatory syndrome in human patients.

Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

45KDa

Swiss-Prot:

Q96BN8

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB,1:500 - 1:2000|IF/ICC,1:50 - 1:200


Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at -20 $^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.


Modification:

Unmodification


DATA:



Western blot analysis of extracts of Rat liver, using OTULIN antibody at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.
br/>Lysates/proteins: 25ug per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
br/>Detection: ECL Basic Kit .
Exposure time: 60s.



Western blot analysis of extracts of MCF7 cells, using OTULIN antibody at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.
br/>Lysates/proteins: 25ug per lane.
br/>Blocking buffer: 3% nonfat dry milk in TBST.
br/>Detection: ECL Basic Kit .
br/>Exposure time: 180s.



Immunofluorescence analysis of NIH/3T3 cells using OTULIN Rabbit

pAb at dilution of 1:50. Blue: DAPI for nuclear staining.

0086-025-68037686 0086-025-68035151

Note:

Tel:

Fax:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com