

# PRODUCT DATA SHEET

Bioworld Technology,Inc.

# MED26 polyclonal antibody

Catalog: BS78041 Host: Rabbit Reactivity: Human, Mouse, Rat

#### **BackGround:**

The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors.

#### **Product:**

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

#### **Molecular Weight:**

70KDa

# **Swiss-Prot:**

O95402

# **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### **Applications:**

WB,1:500 - 1:2000


#### **Storage&Stability:**

Store at  $4\,\mathrm{C}$  short term. Aliquot and store at  $-20\,\mathrm{C}$  long term. Avoid freeze-thaw cycles.


#### **Modification:**

Unmodification

# **DATA:**



Western blot analysis of extracts of various cell lines, using MED26 antibody at 1:1000 dilution.<br/>Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>br/>Lysates/proteins: 25ug per lane.<br/>br/>Blocking buffer: 3% nonfat dry milk in TBST.<br/>Detection: ECL Basic Kit .<br/>br/>Exposure time: 180s.



Western blot analysis of extracts of Raji cells, using MED26 antibody at 1:1000 dilution.<br/>
Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
br/>Lysates/proteins: 25ug per lane.<br/>
Blocking buffer: 3% nonfat dry milk in TBST.<br/>
br/>Detection: ECL Basic Kit.<br/>
Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
br/>Blocking buffer: 3% nonfat dry milk in TBST.<br/>
Blocking buffer: 3% nonfat dry milk in TBST.

#### Note:

For research use only, not for use in diagnostic procedure.

# Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <a href="mailto:info@biogot.com">info@biogot.com</a>
Tel: 0086-025-68037686
Fax: 0086-025-68035151