

# PRODUCT DATA SHEET

Bioworld Technology,Inc.

# [KO Validated] SIRT5 polyclonal antibody

Catalog: BS78689 Host: Rabbit Reactivity: Human, Mouse, Rat

## **BackGround:**

This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class III of the sirtuin family. Alternative splicing of this gene results in multiple transcript variants.

#### **Product:**

1 mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

## **Molecular Weight:**

30KDa

## **Swiss-Prot:**

Q9NXA8

## **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

## **Applications:**

WB,1:500 - 1:2000


# Storage&Stability:

Store at  $4 \,\mathrm{C}$  short term. Aliquot and store at  $-20 \,\mathrm{C}$  long term. Avoid freeze-thaw cycles.


## **Modification:**

Unmodification

## **DATA:**



Western blot analysis of extracts from normal and SIRT5 knockout 293T cells, using SIRT5 antibody at 1:1000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
br/>Lysates/proteins: 25ug per lane.<br/>
br/>Blocking buffer: 3% nonfat dry milk in TBST.<br/>
protection: ECL Basic Kit .<br/>
Exposure time: 1s.



Western blot analysis of extracts of various cell lines, using SIRT5 antibody at 1:1000 dilution.<br/>Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>br/>Lysates/proteins: 25ug per lane.<br/>br/>Blocking buffer: 3% nonfat dry milk in TBST.<br/>Detection: ECL Basic Kit.<br/>br/>Exposure time: 10s.

#### Note:

For research use only, not for use in diagnostic procedure.

## Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u>
Tel: 0086-025-68037686
Fax: 0086-025-68035151