

# PRODUCT DATA SHEET

Bioworld Technology,Inc.

# TAF11 polyclonal antibody

Catalog: BS78955 Host: Rabbit Reactivity: Human, Mouse

## **BackGround:**

Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes a small subunit of TFIID that is present in all TFIID complexes and interacts with TBP. This subunit also interacts with another small subunit, TAF13, to form a heterodimer with a structure similar to the histone core structure. Two transcript variants encoding different isoforms have been found for this gene.

## **Product:**

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

## **Molecular Weight:**

23kDa

# **Swiss-Prot:**

Q15544

# **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum

by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

## **Applications:**

WB,1:500 - 1:2000


# Storage&Stability:

Store at  $4 \,\mathrm{C}$  short term. Aliquot and store at  $-20 \,\mathrm{C}$  long term. Avoid freeze-thaw cycles.

#### **Modification:**

Unmodification

#### **DATA:**



Western blot analysis of extracts of various cell lines, using TAF11 antibody at 1:1000 dilution.<br/>Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>br/>Lysates/proteins: 25ug per lane.<br/>br/>Blocking buffer: 3% nonfat dry milk in TBST.<br/>Detection: ECL Enhanced Kit .<br/>br/>Exposure time: 30s.

### Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <a href="mailto:info@biogot.com">info@biogot.com</a>
Tel: 0086-025-68037686
Fax: 0086-025-68035151