

## CDKN1B/p27KIP1 polyclonal antibody

Catalog: BS79967

Host: Rabbit

Reactivity: Human, Mouse, Rat

#### **BackGround:**

This gene encodes a cyclin-dependent kinase inhibitor, which shares a limited similarity with CDK inhibitor CDKN1A/p21. The encoded protein binds to and prevents the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, and thus controls the cell cycle progression at G1. The degradation of this protein, which is triggered by its CDK dependent phosphorylation and subsequent ubiquitination by SCF complexes, is required for the cellular transition from quiescence to the proliferative state. Mutations in this gene are associated with multiple endocrine neoplasia type IV (MEN4).

#### **Product:**

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

**Molecular Weight:** 

30kDa

**Swiss-Prot:** 

P46527

#### **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

#### **Applications:**

WB,1:500 - 1:2000|IHC,1:50 - 1:100|IF/ICC,1:50 - 1:200


#### Storage&Stability:

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.


#### **Modification:**

Unmodification


**DATA:** 



Western blot analysis of extracts of various cell lines, using CDKN1B/p27KIP1 antibody at 1:3000 dilution.<br/>Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>br/>Lysates/proteins: 25ug per lane.<br/>Blocking buffer: 3% nonfat dry milk in TBST.<br/>br/>Detection: ECL Basic Kit .<br/>br/>Exposure time: 90s.



Immunohistochemistry of paraffin-embedded rat testis using CDKN1B/p27KIP1 antibody at dilution of 1:100 .Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded human breast using CDKN1B/p27KIP1 antibody at dilution of 1:100 .Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

#### Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284


 Fax:
 6122933841

#### Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151



# PRODUCT DATA SHEET

### Bioworld Technology,Inc.



Immunohistochemistry of paraffin-embedded mouse testis using CDKN1B/p27KIP1 antibody at dilution of 1:100 .Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

### Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.		Bioworld technology, co. Ltd.	
Add:	1660 South Highway 100, Suite 500 St. Louis Park,	Add: No 9,	weidi road Qixia District Nanjing, 210046,
	MN 55416,USA.	P. R.	China.
Email:	info@bioworlde.com	Email: info	@biogot.com
Tel:	6123263284	Tel: 0086-	025-68037686
Fax:	6122933841	Fax: 0086-	025-68035151