

CD70 polyclonal antibody

Catalog:	BS80041	Host:	Rabbit	Reactivity: Human, Mouse, Rat
BackGround: The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin sythesis.				The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im- munogen and the purity is > 95% (by SDS-PAGE). Applications: IF/ICC,1:50 - 1:200 Storage&Stability: Store at 4 $^{\circ}$ short term. Aliquot and store at -20 $^{\circ}$ long term. Avoid freeze-thaw cycles. Modification: Unmodification DATA:
Product:				
1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2			0% glycerol,	Immunofluorescence analysis of rat spleen using CD70 Polyclonal An- tibody at dilution of 1:100. Blue: DAPI for nuclear staining.
Molecular Weight:				Note:
Refer to figures Swiss-Prot:				For research use only, not for use in diagnostic procedure.
P32970				
Dentificanting 0 Dentities				

Purification&Purity: