

#### polyclonal antibody **Caspase-1**

Catalog: **BS80060**  Host:

Rabbit

Reactivity: Human, Mouse

# **BackGround:**

This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme. This gene was identified by its ability to proteolytically cleave and activate the inactive precursor of interleukin-1, a cytokine involved in the processes such as inflammation, septic shock, and wound healing. This gene has been shown to induce cell apoptosis and may function in various developmental stages. Studies of a similar gene in mouse suggest a role in the pathogenesis of Huntington disease. Alternative splicing results in transcript variants encoding distinct isoforms.

### **Product:**

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### **Molecular Weight:**

48KDa/20-25KDa

**Swiss-Prot:** 

# P29466

#### **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

#### **Applications:**

WB,1:500 - 1:2000

#### **Storage&Stability:**

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

#### **Modification:**

Unmodification

DATA:

Western blot analysis of extracts of THP-1 cells, using Caspase-1 antibody at 1:500 dilution.THP-1 cells were treated by PMA/TPA at 37°C for overnight and LPS at 37°C for 6 hours.<br/>secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>br/>Lysates/proteins: 25ug per lane.<br/>Blocking buffer: 3% nonfat dry milk in TBST.<br/>br/>Detection: ECL Basic Kit .<br/>Exposure time: 10s.

## Note:

For research use only, not for use in diagnostic procedure.

#### **Bioworld Technology, Inc.** Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 6122933841 Fax:

#### Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. **Email:** info@biogot.com Tel: 0086-025-68037686 0086-025-68035151 Fax: