

COL4A2 polyclonal antibody

Catalog:	BS80513	Host:	Rabbit	Reactivity: Human, Mouse, Rat
BackGround:				munogen and the purity is $> 95\%$ (by SDS-PAGE).
This gene encodes one of the six subunits of type IV col-				Applications:
lagen, the major structural component of basement mem-				WB,1:500 - 1:2000
branes. The C-terminal portion of the protein, known as			in, known as	Storage&Stability:
canstatin, is an inhibitor of angiogenesis and tumor			and tumor	Store at 4 ${}^\circ\!\!{\rm C}$ short term. Aliquot and store at -20 ${}^\circ\!\!{\rm C}$ long
growth. Like the other members of the type IV collagen				term. Avoid freeze-thaw cycles.
gene family, this gene is organized in a head-to-head			head-to-head	Modification:
conformation with another type IV collagen gene so that			gene so that	Unmodification
each gene pair shares a common promoter.				DATA:
Product:				
1mg/ml in PBS with 0.02% sodium azide, 50% glycerol,			0% glycerol,	Western blot analysis of extracts of various cell lines, using COL4A2
pH7.2				antibody at 1:500 dilution. br/>Secondary antibody: HRP Goat An-
Molecular Weight:				ti-Rabbit IgG at 1:10000 dilution. br/>Lysates/proteins: 25ug per
200kDa	200kDa			lane. br/>Blocking buffer: 3% nonfat dry milk in TBST. br/>Detection:
Swiss-Prot:				ECL Basic Kit . br/>Exposure time: 90s.
P08572				Note:
Purification&Purity:				For research use only, not for use in diagnostic procedure.
The antibody was affinity-purified from rabbit antiserum				
by affinity-chromatography using epitope-specific im-				

Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 Fax: 6122933841