

## PRODUCT DATA SHEET

Bioworld Technology,Inc.

# AR polyclonal antibody

Catalog: BS90081 Host: Rabbit Reactivity: Human

### **BackGround:**

Androgens exhibit a wide range of effects on the development, maintenance and regulation of male phenotype and male reproductive physiology. The androgen receptor (AR) is a member of the steroid superfamily of ligand-dependent transcription factors. ARs bind the two biologically active androgens, testosterone (T) and dihydrotestosterone (DHT), with high and nearly identical affinities; however, the rates of association and dissociation of T are about three times more rapid than those of DHT. This difference has resulted in speculation as to whether these differences in binding kinetics could account for the different physiological effects of T and DHT. A striking feature of AR is its rapid degradation in the absence of ligand. It is now well established that androgen binding results in an at least six-fold increase in androgen stability and that ligand-induced stabilization of AR is highly androgen-specific.

### **Product:**

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### **Molecular Weight:**

99 kDa

### **Swiss-Prot:**

P10275(Human)

### **Purification&Purity:**

ProA affinity purified

# **Applications:**

WB:1:500-1:1,000 ICC:1:50-1:200 IHC:1:50-1:100


# Storage&Stability:

Store at +4  $^{\circ}$ C after thawing. Aliquot store at -20  $^{\circ}$ C or -80  $^{\circ}$ C. Avoid repeated freeze / thaw cycles.


### **Specificity:**

AR polyclonal antibody detects endogenous levels of AR protein.

### **DATA:**



ICC staining Androgen receptor in PC-3M cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Androgen receptor in N2A cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

### Note:

For research use only, not for use in diagnostic procedure.

### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u>
Tel: 0086-025-68037686
Fax: 0086-025-68035151