

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



PI3K p110 γ (L89) Peptide

Cat No.: BS9175P

Background

Phosphatidylinositol 3-kinase (PI 3-kinase) is composed of p85 and p110 subunits. p85 lacks PI 3-kinase activity and acts as an adapter, coupling p110 to activated protein tyrosine kinase. Two forms of p85 have been described (p85 α and p85 β), each possessing one SH3 and two SH2 domains. Various p110 isoforms have been identified. p110 α and p110 β interact with p85 α , and p110 α has also been shown to interact with p85 β in vitro. p110 γ expression is restricted to white blood cells. It has been shown to bind p85 α and β , but it apparently does not phosphorylate these subunits. p110 γ seems to have the capacity to autophosphorylate. p110 does not interact with the p85 subunits. It has been shown to be activated by α and β heterotrimeric G proteins.

Swiss-Prot

P48736

Applications

Blocking

Specificity

This peptide can be used with studies using BS9175 PI3K p110 γ (L89) pAb.

Purification & Purity

Synthetic peptide PI3K p110 γ (L89). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.