

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



CD70 (D101) Peptide

Cat No.: BS9259P

Background

The tumor necrosis factor (TNF) receptor family is composed of several type I integral membrane glycoproteins that exhibit homology in their cysteine-rich extracellular domains. Members of this family include FAS, OX40, CD27 and CD30. Ligands for these receptors are often type II transmembrane glycoproteins, as is the case for CD27 and CD30. CD27 is a homodimeric lymphocyte-specific surface antigen present on T and B lymphocytes. Activation of the CD3 complex via the T cell receptor for antigen leads to an increase in CD27 expression. Together, CD27 and its ligand, CD70, generate co-stimulatory signals required for complete T cell activation. CD30 is a surface marker for neoplastic cells of the Hodgkin's lymphoma and related hematologic malignancies. CD30L has been shown to enhance the proliferation of the Hodgkin's cell line HDLM-2, but exerts antiproliferative effects on large cell anaplastic lymphoma cell lines.

Swiss-Prot

P32970

Applications

Blocking

Specificity

This peptide can be used with studies using BS9259 CD70 (D101) pAb.

Purification & Purity

Synthetic peptide CD70 (D101). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.
1660 South Highway 100, Suite 500 St. Louis Park, MN
55416, USA. Email: info@bioworlde.com
Tel: 6123263284 Fax: 6122933841

Bioworld technology, co, Ltd.
No 9, weidi road Qixia District Nanjing, 210046,
P, R.China. Email: info@biogot.com
Tel: +86-025-68037686 Fax: +86-025-68035151