Bioworld Technology CO., Ltd.



GATA-6 (R493) Peptide

Cat No.: **BS9273P**

Background

Members of the GATA family share a conserved zinc finger DNA-binding domain and are capable of binding the WGATAR consensus sequence. GATA-1 is erythroid-specific and is responsible for the regulated transcription of erythroid genes. It is an essential component in the generation of the erythroid lineage. GATA-2 is expressed in embryonic brain and liver, HeLa and endothelial cells, as well as erythroid cells. Studies with a modified GATA consensus sequence, AGATCTTA, have shown that GATA-2 and GATA-3 recognize this mutated consensus while GATA-1 has poor recognition of this sequence. This indicates broader regulatory capabilities of GATA-2 and GATA-3 than GATA-1. GATA-3 is highly expressed in T-lymphocytes. GATA-4, GATA-5 and GATA-6 comprise a subfamily of transcription factors. GATA-4 and GATA-6 are found in heart, pancreas and ovary; lung and liver tissues exhibit GATA-6, but not GATA-4, expression.

Swiss-Prot

Q92908

Applications

Blocking

Specificity

This peptide can be used with studies using BS9273 GATA-6 (R493) pAb.

Purification & Purity

Synthetic peptide GATA-6 (R493). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 $^{\circ}$ C short term. Aliquot and store at -20 $^{\circ}$ C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.