

CDK4 monoclonal antibody

BackGround:Store at 4 °C short term. Aliquot and store at -20Cyclin-dependent kinase activity is regulated by T-loopterm. Avoid freeze-thaw cycles.phosphorylation (Thr172 in the case of CDK4), by theIsotype:	0℃ long
abundance of their cyclin partners, and by association IgG	
with CDK inhibitors of the Cip/Kip or INK family of DATA:	
proteins. Cyclin D-CDK4 complexes are major integra- tors of various mitogenenic and antimitogenic signals.	
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA	
Molecular Weight:	
Calculated MW: 34 kDa; Observed MW: 34 kDa	
Swiss-Prot: Western blot analysis of Cdk4 in K562, Hela lysates using CDI	0K4 anti-
P11802 body.	
Purification&Purity: Note:	
Affinity Purified For research use only, not for use in diagnostic pro	ocedure.
Applications:	
WB: 1/500-1/1000 IF: 1/50-1/200 IP: 1/20	
Storage&Stability:	

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151