

FH monoclonal antibody

Catalog:	MB10919	Host:	Rabbit	Reactivity:	Human,Mouse,Rat,Hamster

BackGround:

Also acts as a tumor suppressor.MiscellaneousThere are 2 substrate-binding sites: the catalytic A site, and the non-catalytic B site that may play a role in the transfer of substrate or product between the active site and the solvent. Alternatively, the B site may bind allosteric effectors .

Product:

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol,

0.01% Sodium azide and 0.05% BSA

Molecular Weight:

Calculated MW: 55 kDa; Observed MW: 49 kDa

Swiss-Prot:

P07954

Purification&Purity:

Affinity Purified

Applications:

WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/20


Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.


Isotype:

IgG

DATA:



Western blot analysis of FH/Fumarase in 3T3, Raw264.7 lysates using FH antibody.



Western blot analysis of FH/Fumarase in Jurkat, rat Brain, C6, CHO-K1 lysates using FH/Fumarase antibody.

Immunohistochemistry analysis of paraffin-embedded Human tonsil us-

ing FH antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Immunocytochemistry analysis of FH in Hela using FH antibody, and DAPI.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151