

NCK2 monoclonal antibody

Catalog:	MB11270	
----------	---------	--

Host: R

```
Rabbit
```

Reactivity: Human, Mouse, Rat

BackGround:

Adapter protein which associates with tyrosine-phosphorylated growth factor receptors or their cellular substrates. Maintains low levels of EIF2S1 phosphorylation by promoting its dephosphorylation by PP1. Plays a role in ELK1-dependent transcriptional activation in response to activated Ras signaling.

Product:

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol,

0.01% Sodium azide and 0.05% BSA

Molecular Weight:

Calculated MW: 43 kDa; Observed MW: 43 kDa

Swiss-Prot:

O43639

Purification&Purity:

Affinity Purified

Applications:

WB: 1/500-1/1000 IF: 1/50-1/200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long

term. Avoid freeze-thaw cycles.

Isotype: IgG

DATA:



Western blot analysis of Nck beta in K562, rat Brain, C6, 3T3, Hela lysates using NCK2 antibody.

Immunocytochemistry analysis of Nck beta in Hela using Nck beta antibody, and DAPI

Note:

For research use only, not for use in diagnostic procedure.